

Minimal Warm Inflation with a heavy QCD axion

Kim V. Berghaus

Burke Institute for Theoretical Physics
California Institute of Technology

Based on: arXiv:2402.13535 (Berghaus, Forslund,
Guevarra)


JCAP03(2020)034 (Berghaus, Graham, Kaplan)



Why Inflation?

- Universe is isotropic
- Universe is flat


$$\frac{\delta T}{T} \approx 10^{-4.5}$$



Planck 2018

Period of accelerated expansion (≈ 60 e-folds) can explain both

Slow-roll Inflation



- Slow-rolling scalar field has negative pressure

$$\ddot{\phi} + 3H\dot{\phi} + V' = 0$$

⊣

Hubble friction


$\delta\phi \sim H$ (quantum fluctuations)

$$\varepsilon_V = \frac{M_{Pl}^2}{2} \left(\frac{V'}{V} \right)^2 \ll 1$$

$$H^2 \approx \frac{V}{3M_{Pl}^2}$$

$$\eta_V = M_{Pl}^2 \frac{V''}{V} \ll 1$$

Slow-roll Inflation


- Inflaton field fluctuations $\delta\phi$ source anisotropies
- Predicts an almost **scale invariant** CMB power spectrum:
- $\Delta_R^2(k) = A_s \left(\frac{k}{k_*}\right)^{n_s-1} \approx -0.035$

 $\sim 10^{-9}$
- $r = 16\varepsilon_V < 0.1$

Challenges

- Simplest models overpredict r
- $\Delta\phi \geq M_{pl}$
- Models undistinguishable

Warm Inflation

Phys.Rev.Lett.75:3218-3221,1995 Berera, 1995



$$T > H$$

$$\rho_{DR} = \frac{\pi^2}{30} g_* T^4$$

- larger non-gaussianities $f_{NL}^{warm} \sim 50$
- Unique bispectral shape

Bastero-Gil, Berera, Moss, Ramos JCAP12(2014)008
Mirbabayi, Gruzinov JCAP02(2023)012

$$3(\Upsilon)\dot{\phi} + V' \approx 0$$

$$4H\rho_{DR} \approx \Upsilon\dot{\phi}^2$$

$$\delta\phi \sim \sqrt{HT} \text{ (classical)}$$

$$\varepsilon_V = \frac{M_{Pl}^2}{2} \frac{3H}{\Upsilon} \left(\frac{V'}{V} \right)^2 \ll 1$$

$$\eta_V = M_{Pl}^2 \frac{3H}{\Upsilon} \frac{V''}{V} \ll 1$$

} suppresses r

} reduces $\Delta\phi$

Microphysics of Warm Inflation

$$\frac{\partial L}{\partial \dot{\phi}} - \frac{d}{dt} \frac{\partial L}{\partial \dot{\phi}} = 0$$

- Couple scalar fields to light degrees of freedom $L_{\text{int}} = -\phi J_{\text{int}}$

$$\ddot{\phi} + 3H\dot{\phi} + V' = -\langle J_{\text{int}} \rangle_{\text{non-eq}}(\phi)$$

$$\langle J_{\text{int}} \rangle_{\text{non-eq}}(\phi) \approx m_{th}^2 \phi + \Upsilon \dot{\phi} + O(\ddot{\phi})$$


Usually $m_{th}^2 \phi \gg \Upsilon \dot{\phi}$, unless symmetry or other reason suppresses m_{th}^2

Minimal Warm Inflation

$$\frac{\partial L}{\partial \dot{\phi}} - \frac{d}{dt} \frac{\partial L}{\partial \ddot{\phi}} = 0$$

- Couple axion to non-Abelian gauge group $L_{\text{int}} = -\phi \frac{\alpha}{16\pi f} \tilde{G}G$ T > H

$$\ddot{\phi} + 3H\dot{\phi} + V' = -\left\langle \frac{\alpha}{16\pi f} \tilde{G}G \right\rangle(\phi)$$




enhanced by sphalerons

$$\left\langle \frac{\alpha}{16\pi f} \tilde{G}G \right\rangle(\phi) \approx m_{th}^2 \cancel{\phi} + \Upsilon \dot{\phi} + O(\ddot{\phi})$$

Not allowed by symmetry

Sphaleron Heating



$$N_{CS}(t) \equiv \int_0^t dt \int dx^3 \text{Tr} \tilde{G} G$$

$$\partial_u K^u = \frac{\alpha}{16\pi f} \tilde{G} G$$

Nonzero $\langle \dot{\phi} \rangle$ biases
sphaleron transitions

$$L_{\text{int}} = -\phi \frac{\alpha}{16\pi f} \tilde{G} G \approx \dot{\phi} K^0$$

Real time sphaleron processes
change topology and efficiently
source particles: $\Upsilon \approx \alpha^5 N^5 \frac{T^3}{f^2}$

Hybrid Warm Inflation


$$V = M_\sigma^{-4} + \frac{1}{2} m_\phi^2 \phi^2; L_{int} = \frac{\alpha}{16\pi f} \tilde{G} G$$

σ drives inflation

ϕ rolls towards ϕ_c

σ reheats into Standard Model

ϕ sources radiation bath




$$T_{RH} \gg T$$

Can we have warm inflation with SM QCD?

$$L_{\text{int}} = -\phi \frac{\alpha_s}{16\pi f} \tilde{G} G \quad \text{SM strong force SU(3)}$$

$$V_{\text{UV}} = \Lambda^4 \left(1 - \cos \frac{\phi}{f_\phi} \right) \quad \text{heavy QCD axion}$$

Sphaleron heating with fermions




- Topology changes built up chiral charge
- Chiral chemical potential counteracts $\langle \dot{\phi} \rangle$

Herranen, arXiv:0906.3136

$$\Upsilon \approx \alpha^5 N^5 \frac{T^3}{f^2}$$

Sphaleron heating with light fermions



Herranen, arXiv:0906.3136

$$\Upsilon \approx \alpha^5 N^5 \frac{T^3}{f^2}$$

QCD sphaleron heating
suppressed due to light
quarks

$$\Upsilon_{eff} = \Upsilon \left(\frac{\Gamma_{chir}}{\Gamma_{chir} + 2f^2/T^2 \Upsilon} \right)$$

$$\Upsilon_{eff} \sim \alpha N \frac{m^2}{f^2} T \text{ if } m \ll \alpha^2 N^2 T$$


Solution: heavy SM quarks
during warm inflation

MWI with a heavy QCD axion

$$V(h) = -\frac{1}{2}\mu^2 h^2 + \lambda(h)h^4$$

The Standard Model quartic of the Higgs runs negative at high energies

Higher dimensional operator can stabilize a second UV Higgs vev



MWI with a heavy QCD axion

$$V(h) = -\frac{1}{2}\mu^2 h^2 + \lambda(h)h^4$$


The Standard Model quartic of the Higgs runs negative at high energies


Higher dimensional operator can stabilize a second UV Higgs vev


QCD quark masses are proportional to v_{UV}

restores

$$\Upsilon_{eff} \rightarrow \Upsilon \text{ if } m \gg \alpha^2 N^2 T$$







Conclusions & Outlook


- Warm inflation is interesting alternative to cold inflation
- Axion coupling to $SU(N)$ is a simple model for thermal particle production via sphaleron heating
- Proof of principle realization for warm inflation with SM particles
- Analysis of chirality violation in full SM sector exciting avenue forward

Thank You

Cold Inflation vs. Minimal Warm Inflation

$$H \ll \Upsilon \propto \alpha^5 \frac{T^3}{f^2}$$

- $\Delta_R^2(k) \propto \delta\phi^2 \propto H^2$
- Tensor to scalar ratio $r \approx 16\varepsilon_V$



$$\varepsilon_V = \frac{M_{Pl}^2}{2} \left(\frac{V'}{V} \right)^2 \ll 1$$


$$\eta_V = M_{Pl}^2 \frac{V''}{V} \ll 1$$

- Small non-gaussianities

- $\Delta_R^2(k) \propto \delta\phi^2 \propto H \textcolor{brown}{T} \left(\frac{\Upsilon}{3H} \right)^{\frac{16}{2}}$

Bastero-Gil, Berera, Ramos *et al* JCAP07(2011)030

- Tensor to scalar ratio $r \approx 0$




$$\varepsilon_V = \frac{M_{Pl}^2}{2} \frac{3H}{\Upsilon} \left(\frac{V'}{V} \right)^2 \ll 1$$

$$\eta_V = M_{Pl}^2 \frac{3H}{\Upsilon} \frac{V''}{V} \ll 1$$

- larger non-gaussianities $f_{NL}^{warm} \sim 50$
- Unique bispectral shape

Bastero-Gil, Berera, Moss, Ramos JCAP12(2014)008

Mirbabayi, Gruzinov JCAP02(2023)012



Reheating

Couple waterfall field to Standard Model: $\frac{\alpha_B \sigma}{16\pi f_B} \tilde{B} B$

